

Bounds on the Parameter of Noncommutativity from Supernova SN1987A

M. Haghghat ¹

Department of Physics Isfahan University of Technology, Isfahan, 84156-83111, Iran

Abstract

We consider supernova SN1987A to find bounds on parameter of noncommutativity, $\theta_{\mu\nu}$. Right handed neutrino in the noncommutative standard model, NCSM, can directly couple to the photon and the Z-gauge boson. Therefore the observed flux of neutrinos from SN1987A can constrain the strength of the new couplings in the NCSM. We obtain two bounds on the NC-parameter, $\Lambda_{NC} = 1/\sqrt{|\theta|}$, with respect to escaping or trapping of the right handed neutrinos inside the supernova which are $\Lambda_{NC} \gtrsim 3.7TeV$ or $\Lambda_{NC} \lesssim 1TeV$, respectively. The excluded region $1TeV \lesssim \Lambda_{NC} \lesssim 3.7TeV$ for the NC-parameter is obtained for the first time. In fact $\Lambda_{NC} \lesssim 1TeV$ is consistent with the existing bounds on Λ_{NC} and raises our hopes to find the NC-effects in the LHC or even in the LEP. Meanwhile $\Lambda_{NC} \gtrsim 3.7TeV$ is more stringent than the other bounds obtained from LEP and LHC considerations. Furthermore, since NC-calculations are perturbative and are correct only up to the energy scale of the NC-system $\lesssim \Lambda_{NC}$, therefore these bounds which are obtained from the energy scale considerably less than the energy scale of LEP and LHC are more reliable.

Noncommutative spaces and their phenomenological aspects have been extensively considered by many authors for a decade [1]-[6]. The natural scale for the noncommutativity effects seems to be the Planck scale which is far from the terrestrial experiment to reach. However, there is no theoretical constraint for the energy scale and one may think this scale might be broken by some mechanism down to the lower energy range and thus be testable by, for example, the Large Hadron Collider (LHC) that is recently started to run. There are two approaches to construct the standard model in the noncommutative space. Nevertheless, for the practical purpose, both versions are based on the perturbative expansion in terms of the parameter of noncommutativity, $\theta_{\mu\nu}$. Actually the region where the θ -expansion is well defined is restricted to $\theta_{\mu\nu}p^\mu q^\nu \lesssim 1$ or $\frac{E}{\Lambda_{NC}} \lesssim 1$ where $E \sim \sqrt{s}$ is the energy of the system. Therefore one can not a priori assume $\Lambda_{NC} \gtrsim \sqrt{s}$ and then compare the new interactions which arise from the perturbative expansion to put bounds on the Λ_{NC} . For instance if $\Lambda_{NC} \sim 200GeV$, the comparison of an obtained amplitude in the NC-space up to the first order of the NC-parameter with the experimental data of the corresponding scattering in $\sqrt{s} \sim 1TeV$ in order to find a bound on Λ_{NC} is obviously incorrect. The existing bounds greater than $\sim 1TeV$ are usually obtained by assuming $\Lambda_{NC} \gtrsim 1TeV$ for the experiments with $\sqrt{s} \gtrsim 1TeV$. Therefore Low-energy experiments tend to provide more reliable limits on $\theta_{\mu\nu}$. There is no doubt that the parameter of noncommutativity is greater than the atomic energy scale. Therefore, one can begin with the atomic constraints such as the Lamb shift in the hydrogen atom [5], the positronium hyperfine splitting or the transitions in the Helium atom [6] to find $\Lambda_{NC} \gtrsim 6GeV, 1MeV, 30GeV$, respectively. Since from the atomic energy scale Λ_{NC} is constrained to be greater than $30GeV$ we can safely apply the perturbative expansion to put bounds on the Λ_{NC} in those processes with the energies up to a few GeV . On the other hand, astrophysical limits can usually provide much more restrictive bounds on strength of new

¹e-mail: mansour@cc.iut.ac.ir

couplings and existence of new particles beyond the standard model than any laboratory experiment could achieve. For example the neutrino burst duration of supernova, SN, 1987A is considered by many authors to find limits on photons escaping into extra dimensions [7], unparticles [8], light dark matter particles [9], neutrino oscillations [10], Kaluza-Klein graviton production [11], sterile-neutrino production [12], axions [13], magnetic moment of the neutrino [14] and the strength of right-handed interactions in the left-right symmetric model[15] among many others. In this letter we consider supernova SN1987A where the energy available for the particles inside the star is about a few hundred of MeV which is far from the confirmed constraint on Λ_{NC} to find new bounds on the parameter of noncommutativity. For this purpose one should consider those new interactions which have not any counterparts in the usual space. In fact the new interactions which appear in the theory as a θ -correction to the existed interactions inside the supernova, can not put a stringent bound on the NC-parameter. As a candidate we can consider the coupling of right handed neutrino with photon in the NCSM which has not any counterpart in the usual standard model. To this end we briefly introduce the direct coupling of the right handed neutrino with photon in the NCSM.

There are two approaches to construct the standard model in the non commutative space. In the simplest one the gauge group is $SU(3)_c \times SU(2)_L \times U(1)_Y$ in which the number of gauge fields, couplings and particles are the same as the ordinary one [3]. In the second approach the gauge group is $U(3) \times U(2) \times U(1)$ which is reduced to $SU(3)_c \times SU(2)_L \times U(1)_Y$ by an appropriate symmetry breaking [2]. However, in the both versions, new interactions with respect to the ordinary standard model will appear due to the star product and the Seiberg-Witten, SW, map. Here we consider the former case and denote an infinitesimal non-commutative local gauge transformation of the fields contents of the theory as $\delta\hat{\Psi} = i\rho_{\Psi}(\Lambda) \star \hat{\Psi}$, where Λ is a gauge parameter and $\rho_{\Psi}(\Lambda)$ is a representation which is carried by the matter or Higgs fields. The fields with hat are the non-commutative fields which can be written as a function of ordinary fields using appropriate SW-maps. The right handed neutrino, $\hat{\nu}_R$, as a neutral-hyper-charged particle in the standard model can be transformed as $\delta\hat{\nu}_R = i\hat{\Lambda}' \star \hat{\nu}_R - i\hat{\nu}_R \star \hat{\Lambda}'$ to preserve the gauge symmetry. In fact in a non-commutative setting the non-commutative gauge boson B_{μ} , compatible with the non-commutative gauge transformation, couples to a neutral matter field $\hat{\Psi}^0$ as $\hat{D}_{\mu}\hat{\Psi}^0 = \partial_{\mu}\hat{\Psi}^0 - i[\hat{B}_{\mu}, \hat{\Psi}^0]$, with $\delta\hat{\Psi}^0 = i[\hat{\Lambda} \star, \hat{\Psi}^0]$. In other words, to introduce the neutrino-photon interaction in the minimal NCSM, one can define the adjoint representation in the covariant derivative for the neutral particle as is already done in the NCQED. In other words in the frame work of NCQED it was shown that the neutral particles interact with photons if they transform under $U(1)$ in a similar way as in the adjoint representation of a non-Abelian gauge theory [16]. The main difference in the NCSM is $U(1)_Y$ instead of $U(1)_{EM}$. Therefore, in a gauge invariant manner, only neutral hyper charge particle in the standard model can couple to the hyper gauge field. The only particle with zero hyper charge in the SM is the right handed neutrino therefore the covariant derivative for this particle, to the lowest order, can be written as $\hat{D}_{\mu}\hat{\psi}_{\nu_R} = \partial_{\mu}\hat{\psi}_{\nu_R} + e\theta^{\nu\rho}\partial_{\nu}\hat{B}_{\mu}\partial_{\rho}\hat{\psi}_{\nu_R}$, [17], in which $\hat{\psi}_{\nu_R}$ and $\hat{B}$, respectively, denote the NC-fields of the right handed neutrino and the hyper charge with their own expansion up to the lowest order in the NC-space as $\hat{\psi}_{\nu_R} = \psi_{\nu_R} + e\theta^{\nu\rho}B_{\rho}\partial_{\nu}\psi_{\nu_R}$ and $\hat{B}_{\mu} = B_{\mu} + e\theta^{\nu\rho}B_{\rho}[\partial_{\nu}B_{\mu} - \frac{1}{2}\partial_{\mu}B_{\nu}]$. Consequently, Lagrangian density for the right handed neutrino part of NCSM can be written as follows [17]

$$\begin{aligned}\mathcal{L}_{\nu_R} &= i\bar{\psi}\not{\partial}\psi + ie\theta^{\mu\nu}[\partial_{\mu}\bar{\psi}B_{\nu}\gamma^{\rho}(\partial_{\rho}\psi) \\ &\quad - \partial_{\rho}\bar{\psi}B_{\nu}\gamma^{\rho}(\partial_{\mu}\psi) + \bar{\psi}(\partial_{\mu}B_{\rho})\gamma^{\rho}(\partial_{\nu}\psi)],\end{aligned}\tag{1}$$

where B in terms of the photon and the Z -gauge boson fields is $B = \cos\theta_W A - \sin\theta_W Z$. Therefore

the Feynman rules for $\gamma\nu\bar{\nu}$ and $Z\nu\bar{\nu}$ vertices can be obtained from the Lagrangian (1) as:

$$\Gamma_{\gamma\nu\bar{\nu}}^{\mu} = i\frac{e}{2}\cos\theta_W(1 + \gamma_5)(\theta^{\mu\nu}k_{\nu}\not{q} + \theta^{\rho\mu}q_{\rho}\not{k} + \theta^{\nu\rho}k_{\nu}q_{\rho}\gamma^{\mu}), \quad (2)$$

and

$$\Gamma_{Z\nu\bar{\nu}}^{\mu} = -i\frac{e}{2}\sin\theta_W(1 + \gamma_5)(\theta^{\mu\nu}k_{\nu}\not{q} + \theta^{\rho\mu}q_{\rho}\not{k} + \theta^{\nu\rho}k_{\nu}q_{\rho}\gamma^{\mu}). \quad (3)$$

Since the other particles in the SM, even the left handed neutrino, all have nonzero hyper charge, the remaining parts of the SM in the noncommutative space do not change. It should be noted that for neutrino as a neutral particle in the NCQED, as well as QED, in contrast with the standard model, there is no constraint on the mass or even the chirality of the neutrino. In the standard model, neutrino is massless and only the left handed one has weak interaction while the right handed neutrino, if existing, has an expectator role in all reactions.

Now we ready to calculate the impact of this new interaction on the luminosity of the supernova 1987A. Inside a supernova, several particle and nucleon processes contribute to neutrino production, absorption and neutrino scattering. In fact the neutrino interactions are essential in producing and powering the supernova explosion. The neutrino spectrum is always characterized by a short peak of electron neutrinos generated during the neutronization phase carrying 1% of the total energy and a second release of neutrinos of all flavors stemming from thermal pair processes (pair annihilation, plasmon decay, photoneutrino and Bremsstrahlung) releasing 99% of the energy. Meanwhile since the nucleon density is huge, the dominant lowest-order process for the emission of new particles in a supernova core tends to be nucleon bremsstrahlung. However in the calculation of nucleon-nucleon interactions, one encounters the nonperturbative part of the strong interactions which is not well known. Furthermore, if the interaction of new particles with all nucleons is the same, dipole emission in the nucleon bremsstrahlung is suppressed in the nonrelativistic limit. Therefore, one expects that the smaller leptonic density be compensated by the fact that the pair annihilation channel is not suppressed [18]-[19]. Consequently pair annihilation can be considered as a competent source of new particles producing in the core of supernova. In the absence of the left-right symmetric model for the electroweak interaction, the right handed neutrinos can be produced in the supernova via pair annihilation in the noncommutative space

$$f + \bar{f} \rightarrow \nu_R + \bar{\nu}_R. \quad (4)$$

As one can easily see, although at the lowest order in the NC-space, the annihilation can be proceeded via the one photon exchange as well as the one Z-boson exchange, the latter one is highly suppressed because of the Z-boson propagator which leads to the additional power of the Fermi coupling constant. Therefore we only consider the one photon exchange diagram at the tree level and its crossed amplitude for charged fermions specially for the electrons and muons. Indeed the temperature of the proto-neutron star is sufficient to generate a muonic number density as large as the number density of the massless fermions. Once the right handed neutrinos are produced, they have mean-free-paths in the supernova core which are determined via the cross sections for the scattering processes as follows

$$\begin{aligned} \nu_R + f &\rightarrow \nu_R + f, \\ \nu_R + p &\rightarrow \nu_R + p. \end{aligned} \quad (5)$$

In light of this situation, here we consider the observed neutrinos from SN 1987A [20] which is in agreement with the theoretical picture for the Type-II supernovae. The impact of the new interaction on the data of the SN 1987A is twofold:

1. If the strength of the new coupling which depends inversely on the parameter of noncommutativity, Λ_{NC} , leads to a mean free path larger than the core radius, the right handed neutrinos can escape the proto-neutron star. This provides a new channel for energy loss which increases with decreasing Λ_{NC} . Therefore the comparison of the ν_R -luminosity with the well known data from the SN 1987A can put a lower bound on Λ_{NC} .
2. Meanwhile the absorption cross section increases as Λ_{NC} decreases and therefore for a certain range of Λ_{NC} the neutron star becomes optically thick for the right handed neutrinos. In this case they have mean free path smaller than the core radius and can be trapped inside the ν_R -sphere. In this situation the luminosity can be approximated by black body radiation which in turn decreases as Λ_{NC} decreases. In other words, there is an upper bound on the parameter of noncommutativity in this case.

Free Streaming Case: In the NC-space there is no constraint on mass of the right handed neutrino. Therefore in the NC-space, the light right handed neutrinos can be produced in the SN via the fermion pair annihilation given in Eq.(4). By using the $\gamma\nu_R\nu_R$ -vertex given in Eq.(2) and neglecting the mass of neutrino, the spin-averaged amplitude in terms of the Mandelstam variables can be obtained as

$$|\overline{\mathcal{M}}|^2 = e^4 \cos^4 \theta_W [k \cdot \theta \cdot p]^2 \frac{t^2 + u^2 + m_f^2 s}{s^2}, \quad (6)$$

where k and p are the momenta of f and ν_R , respectively. Here for simplicity we calculate the total cross section in the center of mass frame. To this end we define

$$\begin{aligned} \vec{\theta}_s &= (\theta_{23}, \theta_{31}, \theta_{12}) = (\theta_{s\parallel}, \theta_{s\perp}), \\ \vec{\theta}_t &= (\theta_{01}, \theta_{02}, \theta_{03}) = (\theta_{t\parallel}, \theta_{t\perp}), \end{aligned} \quad (7)$$

where $\perp$ and $\parallel$ mean perpendicular to and parallel with the surface contains $\vec{p}$ and $\vec{k}$. For the electron pair annihilation, we can neglect the mass of electron with respect to the center of mass energy, and after averaging on the angle of $\vec{\theta}$ with the $\vec{p}$ and $\vec{k}$, to find

$$\sigma(ee^+ \rightarrow \nu_R \bar{\nu}_R) \simeq \frac{e^4 \cos^4 \theta_W}{30\pi} \Theta^2 E^2 = 0.01 \left(\frac{M_W}{\Lambda_{NC}} \right)^4 G_F^2 E^2, \quad (8)$$

where $\Theta = [5 |\theta_{t\parallel}|^2 / 8 + 3 |\theta_{s\perp}|^2 / 8]$ and $\Lambda_{NC} = 1/\sqrt{\Theta}$.

If the volume energy-loss rate of the SN by any new mechanism becomes greater than $Q \sim 3 \times 10^{33} \text{ ergs cm}^{-3} \text{ s}^{-1}$ then it will remove sufficient energy from the explosion to invalidate the current understanding of the SN1987A-neutrino signal [21]. Meanwhile the new mechanism in the NC-space, Eq.(4), leads to the energy emitted by the supernova per unit time and unit volume as follows

$$Q = n(e^+)n(e^-)\langle \sigma(ee^+ \rightarrow \nu_R \bar{\nu}_R) v_{rel} (E_\nu + E_{\bar{\nu}}) \rangle, \quad (9)$$

where $n(e^\pm)$ is the number density of electron (positron) and v_{rel} is the relative velocity of e^- and e^+ . Therefore Eq.(8) and Eq.(9) for zero chemical potential and equal number density of electron and neutron result in

$$Q = 2.5 \times 10^{-3} \left(\frac{M_W}{\Lambda_{NC}} \right)^4 G_F^2 \frac{16}{3\pi^4} T^9 \int_0^\infty \frac{x^3}{e^x + 1} dx \int_0^\infty \frac{x^4}{e^x + 1} dx, \quad (10)$$

where for $T = 50$ one finds $Q \sim (\frac{M_W}{\Lambda_{NC}})^4 \times 10^{40} \text{ ergs cm}^{-3} \text{ s}^{-1}$ or $\Lambda_{NC} \gtrsim 4.5 \text{ TeV}$. The presence of a chemical potential reduces the number of pairs but for the muon with the rest mass of 106 MeV ignoring the chemical potential does not lead to a substantial error. However for the pairs of muons, there is a suppression factor with respect to the electrons of about 0.5 coming from the relative velocity of muons. Therefore for $T = 50$ one finds $\Lambda_{NC} \gtrsim 3.7 \text{ TeV}$.

Trapping Case: If the interaction of the right handed neutrino in the NC-space is strong enough it can be trapped and its subsequent thermalization can reduce the luminosity. The most relevant reactions which contribute to trapping are given in Eq.(5). In order to find the luminosity in this case we need to estimate the radius of the ν_R -sphere. Now for simplicity we neglect the mass of the charged fermion and restrict ourselves to the space noncommutativity (i.e. $\theta_t = 0$), to obtain the differential cross section as follows

$$\frac{d\sigma}{d\Omega} = \frac{e^4 \cos^4 \theta_W}{256\pi^2} E^2 |\theta_{s\perp}|^2 \frac{(1 + \cos\alpha)(4 + (1 + \cos\alpha)^2)}{1 - \cos\alpha}, \quad (11)$$

where α is the scattering angle. It should be noted that the differential cross section as usual has a singularity at $\cos\alpha = 1$ but here the singularity is only logarithmic. To obtain the total cross section we need a cutoff to regularize the divergence when α goes to zero. In fact neutrinos are confined in the ν_R -sphere with a radius of the order of the core radius, R_C , that constrains their transverse momentum to be at least $p_\perp \sim \frac{1}{R_C}$ or the scattering angle should be greater than $\alpha = \frac{p_\perp}{E} \sim 10^{-19}$. Alternatively, the screening effect in the core can provide a mass of the order of $\mu = V^{-\frac{1}{3}}$ to the photon which in turn changes the variable $t \rightarrow t - \mu^2$ or $1 - \cos\alpha$ to $1 - \cos\alpha + \frac{\mu^2}{2E^2}$ which is regular at $\cos\alpha = 1$ and is equivalent to $\alpha \gtrsim 10^{-19}$. Therefore the total cross section can be easily obtained as

$$\sigma = 3.28 \left(\frac{M_W}{\Lambda_{NC}}\right)^4 G_F^2 E^2. \quad (12)$$

Eq.(12) can be used to calculate the mean free path of the right handed neutrino. For constant temperature and core number density one can easily find the right handed neutrino mean free path as

$$l_{\nu_R} = \frac{1}{n\sigma} = \{95.12 \left(\frac{M_W}{\Lambda_{NC}}\right)^4 \left(\frac{n}{n_C}\right) \left(\frac{T}{50}\right)^2\}^{-1} m, \quad (13)$$

where $E^2 \sim 10T^2$ and $n_C = 2.4 \times 10^{44} \text{ m}^{-3}$. In the trapping case $l_{\nu_R} \lesssim R_C$ which for $R_C \sim 10^4 \text{ m}$ and $T = 50 \text{ MeV}$ leads to $\Lambda_{NC} \lesssim 31 M_W$. If one considers the variation of temperature and number density in the core a more realistic bound on the Λ_{NC} can be found. For instance, if $n(R) = n_C \left(\frac{R}{R_C}\right)^m$ and $T(R) = T_C \left(\frac{R}{R_C}\right)^{\frac{m}{3}}$, for an appropriate value of m see [12] and [15], one finds from Eq.(12) $\Lambda_{NC} \lesssim 1.1 \text{ TeV}$.

Now let us summarize our results. In this letter we showed that the direct coupling of the right handed neutrino with photon in the NCSM can provide a new channel for the energy loss of SN 1987A. Since NC-calculations are perturbative and are correct only up to the energy scale of the NC-system $\lesssim \Lambda_{NC}$, therefore SN 1987A as a system with energy scale of a few hundred of MeV is an excellent candidate to find bounds on Λ_{NC} . We found for the weak coupling that $\Lambda_{NC} \gtrsim 3.7 \text{ TeV}$ which is more stringent than the other bounds obtained from LEP and LHC considerations. Meanwhile for the strong coupling, the trapping case, we obtained $\Lambda_{NC} \lesssim 1 \text{ TeV}$. In fact constraints on the energy loss of SN 1987A exclude the range $1 \text{ TeV} \lesssim \Lambda_{NC} \lesssim 3.7 \text{ TeV}$ for the NC-parameter. The lower bound can be considered seriously because the obtained bounds from LEP and LHC considerations are usually

based on the fact that $\Lambda_{NC} \gtrsim$ the energy scale of the system which may not be correct. Therefore $\Lambda_{NC} \lesssim 1TeV$ is consistent with the existing bounds on Λ_{NC} and raises our hopes to find the NC-effects in the LHC or even in the LEP. Finally it should be noted that the well known processes which can successfully describe the energy loss of SN 1987A in the ordinary space also receive the NC-corrections. But these corrections to the usual vertices are too small to change the energy loss of the supernova. For instance, in the production of the left handed neutrino in the ordinary space the cross section is of the order of $\sigma_0 \sim G^2 E^2$ and at the lowest order in the NC-space its correction is of the order of $\sigma_{NC} \sim \theta G^2 E^4$ where for $\Lambda = \frac{1}{\sqrt{\theta}} \sim 1TeV$ and for the energy range $\sim 10MeV$ the fraction $\frac{\sigma_{NC}}{\sigma_0} \sim \theta E^2 \sim 10^{-10}$ is negligible. Therefore the energy loss of the supernova via the usual channels do not change in the NC-space.

References

- [1] N. Seiberg and E. Witten, JHEP **9909**, 032 (1999).
- [2] M. Chaichian, P. Prešnajder, M. M. Sheikh-Jabbari and A. Tureanu, Phys. Lett. B **526**, 132 (2002); M. Chaichian, P. Prešnajder, M. M. Sheikh-Jabbari and A. Tureanu, Eur. Phys. J. **C29**, 413 (2003).
- [3] X. Calmet, B. Jurčo, P. Schupp, J. Wess and M. Wohlgenannt, Eur. Phys. J. **C23**, 363 (2002); B. Melić, K. Passek-Kumerički, J. Trampetić, P. Schupp and M. Wohlgenannt, Eur. Phys. J. **C42**, 483(2005).
- [4] M. Hayakawa, Phys. Lett. B **478**, 394 (2000); M. Haghghat and F. Loran, Mod. Phys. Lett. A **16**, 1435 (2001); I. Hinchliffe, N. Kersting, and Y. L. Ma, Int. J. Mod. Phys. A **19**, 179 (2004); M. Haghghat and M. M. Ettefaghi, Phys. Rev. D **70**, 034017 (2004); A. Alboteanu, T. Ohl and R. Rckl, Phys. Rev. D **74**, 096004 (2006); M. Mohammadi Najafabadi, Phys. Rev. D **74**, 025021 (2006); M. M. Ettefaghi and M. Haghghat, Phys. Rev. D **75**, 125002 (2007); C. P. Martin, D. Sanchez-Ruiz and C. Tamarit, JHEP **0702**, 065 (2007).
- [5] M. Chaichian, M. M. Sheikh- Jabbari, and A. Tureanu, Phys. Rev. Lett. **86**, 2716 (2001); A. Stern, Phys. Rev. Lett. **100**, 061601 (2008).
- [6] M. Haghghat, S. M. Zebarjad and F. Loran, Phys. Rev. D. **66** , 016005 (2002); M. Haghghat and F. Loran, Phys. Rev. D. **67**, 096003 (2003).
- [7] A. Friedland and M. Giannotti, Phys. Rev. Lett. **100**, 031602 (2008).
- [8] H. Davoudiasl, Phys. Rev. Lett. **99**, 141301 (2007).
- [9] P. Fayet, D. Hooper, and G. Sigl, Phys. Rev. Lett. **96**, 211302 (2006).
- [10] T. Yoshida, T. Kajino, H. Yokomakura, K. Kimura, A. Takamura, and Dieter H. Hartmann, Phys. Rev. Lett. **96**, 091101 (2006).
- [11] S. Hannestad and G. G. Raffelt, Phys. Rev. Lett. **87**, 051301 (2001).
- [12] E. W. Kolb, R. N. Mohapatra, and V. L. Teplitz, Phys. Rev. Lett. **77**, 3066 (1996).

- [13] G. Raffelt and D. Seckel, Phys. Rev. Lett. **67**, 2605 (1991).
- [14] R. Barbieri and R. N. Mohapatra, Phys. Rev. Lett. **61**, 27 (1988).
- [15] G. Raffelt and D. Seckel, Phys. Rev. Lett. **60**, 1793(1988); R. Barbieri and R. N. Mohapatra, Phys. Rev. D. **39**, 1229(1989).
- [16] P. Schupp, J. Trampetic, J. Wess and G. Raffelt, Eur. Phys. J. C **36**, 405 (2004); H. Grosse and Y. Liao, Phys. Rev. D **64**, 115007 (2001); H. Grosse and Y. Liao, Phys. Lett. B **520**, 63 (2001).
- [17] M. Haghigat, M. M. Ettefaghi and M. Zeinali, Phys. Rev. D **73**, 013007 (2006); M. M. Ettefaghi and M. Haghigat, **77**, 056009 (2008).
- [18] R. Buras, H. T. Janka, M. T. Keil, G. G. Raffelt and M. Rampp, Astrophys. J. **587**, 320 (2003).
- [19] S. Hannestad, G. G. Raffelt, and Yvonne Y. Y. Wong, arXiv:0708.1404.
- [20] K. Hirata et al., Phys. Rev. Lett. **58**, 1490 (1987) ; R. M. Bionta et al., Phys. Rev. Lett. **58**, 1494 (1987).
- [21] G. G. Raffelt, Stars as Laboratories for Fundamental Physics, (Chicago University Press) (1996).